

6/9/05

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1612RXD

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page URLs for STN Seminar Schedule - N. America
NEWS	2		"Ask CAS" for self-help around the clock
NEWS	3	FEB 25	CA/CAPLUS - Russian Agency for Patents and Trademarks (ROSPATENT) added to list of core patent offices covered
NEWS	4	FEB 28	PATDPAFULL - New display fields provide for legal status data from INPADOC
NEWS	5	FEB 28	BABS - Current-awareness alerts (SDIs) available
NEWS	6	FEB 28	MEDLINE/LMEDLINE reloaded
NEWS	7	MAR 02	GBFULL: New full-text patent database on STN
NEWS	8	MAR 03	REGISTRY/ZREGISTRY - Sequence annotations enhanced
NEWS	9	MAR 03	MEDLINE file segment of TOXCENTER reloaded
NEWS	10	MAR 22	KOREAPAT now updated monthly; patent information enhanced
NEWS	11	MAR 22	Original IDE display format returns to REGISTRY/ZREGISTRY
NEWS	12	MAR 22	PATDPASPC - New patent database available
NEWS	13	MAR 22	REGISTRY/ZREGISTRY enhanced with experimental property tags
NEWS	14	APR 04	EPFULL enhanced with additional patent information and new fields
NEWS	15	APR 04	EMBASE - Database reloaded and enhanced
NEWS	16	APR 18	New CAS Information Use Policies available online
NEWS	17	APR 25	Patent searching, including current-awareness alerts (SDIs), based on application date in CA/CAPLUS and USPATFULL/USPAT2 may be affected by a change in filing date for U.S. applications.
NEWS	18	APR 28	Improved searching of U.S. Patent Classifications for U.S. patent records in CA/CAPLUS
NEWS	19	MAY 23	GBFULL enhanced with patent drawing images
NEWS	20	MAY 23	REGISTRY has been enhanced with source information from CHEMCATS
NEWS	21	MAY 26	STN User Update to be held June 6 and June 7 at the SLA 2005 Annual Conference
NEWS	22	JUN 06	STN Patent Forums to be held in June 2005
NEWS	23	JUN 06	The Analysis Edition of STN Express with Discover! (Version 8.0 for Windows) now available
NEWS EXPRESS			JANUARY 10 CURRENT WINDOWS VERSION IS V7.01a, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 10 JANUARY 2005
NEWS HOURS			STN Operating Hours Plus Help Desk Availability
NEWS INTER			General Internet Information
NEWS LOGIN			Welcome Banner and News Items
NEWS PHONE			Direct Dial and Telecommunication Network Access to STN
NEWS WWW			CAS World Wide Web Site (general information)

10714255

6/9/05

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 18:31:34 ON 08 JUN 2005

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'CAPLUS' ENTERED AT 18:31:48 ON 08 JUN 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 8 Jun 2005 VOL 142 ISS 24

FILE LAST UPDATED: 7 Jun 2005 (20050607/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s (Becker and 1988 and 5 and 12 and 1053)/so

428 BECKER/SO

448999 1988/SO

2410080 5/SO

1062946 12/SO

9339 1053/SO

L1 0 (BECKER AND 1988 AND 5 AND 12 AND 1053)/SO

=> s (1988 and 5 and 12 and 1053)/so

448999 1988/SO

2410080 5/SO

1062946 12/SO

9339 1053/SO

L2 1 (1988 AND 5 AND 12 AND 1053)/SO

=> d abs bib hitstr


10714255

6/9/05

10714255

6/9/05

L2 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2005 ACS on STN
G1



AB Nil [H₂L = I (R = H, Me, Et, iso-Pr, Bu; R₁ = CPhCPh, CH₂CH₂OMe)] were prepared from NiCl₂, dioxime and H₃BO₃ in ROH. All the compds. are spin-paired, crystalline intensely colored solids. IR and electronic spectra show the complexes are isostructural and nearly square planar.

AN 1989:608105 CAPLUS

DN 111:208105

TI Alkoxy boron-bridged macrocyclic compounds of nickel(II) with 1,1',8,8'-tetraalkoxy-1,8-dibora-4,5,11,12-tetraphenyl-3,6,10,13-tetraaza-2,7,9,14-tetraoxatetradecacyclo-3,5,10,12-tetraene and 1,1',9,9'-tetraalkoxy-1,9-dibora-4,6,12,14-tetramethyl-3,7,11,15-tetraaza-2,8,10,16-tetraoxacyclohexadeca-3,6,11,14-tetraene

AU Rai, H. C.; Tiwary, Jaya; Kumari, Rekha

CS L S College, Univ. Bihar, Muzaffarpur, India

SO Indian Journal of Chemistry, Section A: Inorganic, Physical, Theoretical & Analytical (1988), 27A(12), 1053-5

CODEN: IJCA DU; ISSN: 0376-4710

DT Journal

LA English

10714255

6/9/05

=> s (1998 and 5 and 12 and 1053)/so

712861 1998/SO

2410080 5/SO

1062946 12/SO

9339 1053/SO

L3 2 (1998 AND 5 AND 12 AND 1053)/SO

=> d abs bib hitstr 1-2

6/9/05

L3 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2005 ACS on STN
AB Eubacterial proteins are synthesized with a formyl group at the N-terminus which is hydrolytically removed from the nascent chain by the mononuclear Fe-containing enzyme, peptide deformylase (1). The catalytic efficiency strongly depends on the identity of the bound metal. Here, the authors determined by x-ray crystallog. the Fe2+, Ni2+, and Zn2+ forms of Escherichia coli I and a crystal structure of I complexed with the reaction product, Met-Ala-Ser. The structure of the I complex, with the tripeptide bound at the active site, suggested detailed models for the mechanism of substrate recognition and catalysis. Differences in the protein structures due to the identity of the bound metal were extremely small and accounted only for the observation that Zn2+ bound more tightly than Fe2+ or Ni2+. The striking loss of catalytic activity of the Zn2+ form could be caused by its reluctance to change between tetrahedral and 5-fold metal coordination believed to occur during catalysis.
AN 1998:797694 CAPLUS
DN 130:150294
TI Iron center, substrate recognition and mechanism of peptide deformylase
AU Becker, Andreas; Schlichting, Ilme; Kabsch, Wolfgang; Groche, Dieter; Schultz, Sabine; Wagner, A. F. Volker
CS Max-Planck-Institut für medizinische Forschung, Abteilung Biophysik, Heidelberg, 69120, Germany
SO Nature Structural Biology (1998), 5(12), 1053-1058
CODEN: NSBIEW; ISSN: 1072-8368
PB Nature America
DT Journal
LA English
RE.CNT 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2005 ACS on STN
AB The objective of this investigation was to determine if the decrease of cytochrome P 4502E1 (CYP2E1) in hepatomas is related to the tumor growth rate. There was a significant correlation between N-nitrosodimethylamine (NDMA) demethylase activities and CYP2E1 protein levels in rat liver and hepatomas. The levels of NDMA demethylase activities and CYP2E1 protein content were lower in hepatomas than in the corresponding host livers. NDMA demethylase activities and CYP2E1 protein levels were greater in hepatomas of slow and intermediate growth rate than in fast growing hepatomas. A similar trend was also observed with CYP2E1 mRNA levels. The results demonstrated an inverse relation between growth rate of rat hepatomas and the expression of CYP2E1.
AN 1998:307705 CAPLUS
DN 129:79913
TI Differential expression of cytochrome P4502E1 (CYP2E1) in Morris hepatomas and livers of tumor bearing rats
AU Hu, Jennifer J.; Hong, Jun-Yan; Lea, Michael A.; Yang, Chung S.
CS Laboratory Cancer Research, Dep. Chemical Biology Pharmacognosy, College Pharmacy, Rutgers Univ., Piscataway, NJ, 08855, USA
SO International Journal of Oncology (1998), 12(5), 1049-1053
CODEN: IJONES; ISSN: 1019-6439
PB International Journal of Oncology
DT Journal
LA English
RE.CNT 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

10714255

6/9/05

=> FIL STNGUIDE

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	34.32	34.53

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-2.19	-2.19

FILE 'STNGUIDE' ENTERED AT 18:34:17 ON 08 JUN 2005
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE
AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.
LAST RELOADED: Jun 3, 2005 (20050603/UP).

=> logoff y

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.30	34.83

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	0.00	-2.19

STN INTERNATIONAL LOGOFF AT 18:37:11 ON 08 JUN 2005

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1612RXD

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1	Web Page URLs for STN Seminar Schedule - N. America
NEWS	2	"Ask CAS" for self-help around the clock
NEWS	3 FEB 25	CA/CAPLUS - Russian Agency for Patents and Trademarks (ROSPATENT) added to list of core patent offices covered
NEWS	4 FEB 28	PATDPAFULL - New display fields provide for legal status data from INPADOC
NEWS	5 FEB 28	BABS - Current-awareness alerts (SDIs) available
NEWS	6 FEB 28	MEDLINE/LMEDLINE reloaded
NEWS	7 MAR 02	GBFULL: New full-text patent database on STN
NEWS	8 MAR 03	REGISTRY/ZREGISTRY - Sequence annotations enhanced
NEWS	9 MAR 03	MEDLINE file segment of TOXCENTER reloaded.

10714255

6/9/05

NEWS 10 MAR 22 KOREAPAT now updated monthly; patent information enhanced
NEWS 11 MAR 22 Original IDE display format returns to REGISTRY/ZREGISTRY
NEWS 12 MAR 22 PATDPASPC - New patent database available
NEWS 13 MAR 22 REGISTRY/ZREGISTRY enhanced with experimental property tags
NEWS 14 APR 04 EPFULL enhanced with additional patent information and new fields
NEWS 15 APR 04 EMBASE - Database reloaded and enhanced
NEWS 16 APR 18 New CAS Information Use Policies available online
NEWS 17 APR 25 Patent searching, including current-awareness alerts (SDIs), based on application date in CA/CAPLUS and USPATFULL/USPAT2 may be affected by a change in filing date for U.S. applications.
NEWS 18 APR 28 Improved searching of U.S. Patent Classifications for U.S. patent records in CA/CAPLUS
NEWS 19 MAY 23 GBFULL enhanced with patent drawing images
NEWS 20 MAY 23 REGISTRY has been enhanced with source information from CHEMCATS
NEWS 21 MAY 26 STN User Update to be held June 6 and June 7 at the SLA.2005 Annual Conference
NEWS 22 JUN 06 STN Patent Forums to be held in June 2005
NEWS 23 JUN 06 The Analysis Edition of STN Express with Discover! (Version 8.0 for Windows) now available

NEWS EXPRESS JANUARY 10 CURRENT WINDOWS VERSION IS V7.01a, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 10 JANUARY 2005

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 11:58:13 ON 09 JUN 2005

=> file registry

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'REGISTRY' ENTERED AT 11:58:21 ON 09 JUN 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 American Chemical Society (ACS)

10714255

6/9/05

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 8 JUN 2005 HIGHEST RN 851931-88-9
DICTIONARY FILE UPDATES: 8 JUN 2005 HIGHEST RN 851931-88-9

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 18, 2005

Please note that search-term pricing does apply when conducting SmartSELECT searches.


```
*****
*
* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *
*
*****
```

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10714255.str



chain nodes :

1 2 3 4 17 24 25 26 27 28 29 30 31

ring nodes :

5 6 7 8 9 10 11 12 13 14 15 16 18 19 20 21 22 23

chain bonds :

1-2 1-3 1-4 1-24 3-5 4-11 13-17 17-18 24-25 25-26 26-27 26-28 28-29
29-30 30-31

ring bonds :

10714255

6/9/05

5-6 5-10 6-7 7-8 8-9 9-10 11-12 11-16 12-13 13-14 14-15 15-16 18-19
18-23 19-20 20-21 21-22 22-23
exact/norm bonds :
1-2 1-3 1-4 3-5 4-11 13-17 17-18 24-25 25-26 26-27 28-29 29-30 30-31
exact bonds :
1-24 26-28
normalized bonds :
5-6 5-10 6-7 7-8 8-9 9-10 11-12 11-16 12-13 13-14 14-15 15-16 18-19
18-23 19-20 20-21 21-22 22-23

Match level :

1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:CLASS 18:Atom 19:Atom
20:Atom 21:Atom 22:Atom 23:Atom 24:CLASS 25:CLASS 26:CLASS 27:CLASS
28:CLASS 29:CLASS 30:CLASS 31:CLASS

L1 STRUCTURE UPLOADED

=> d l1

L1 HAS NO ANSWERS

L1 STR

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

Structure attributes must be viewed using STN Express query preparation.

=> s l1

SAMPLE SEARCH INITIATED 11:58:40 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 0 TO ITERATE

100.0% PROCESSED 0 ITERATIONS

0 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**

BATCH **COMPLETE**

PROJECTED ITERATIONS: 0 TO 0

PROJECTED ANSWERS: 0 TO 0

L2 0 SEA SSS SAM L1

=> s l1 ful

FULL SEARCH INITIATED 11:58:53 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 4 TO ITERATE

100.0% PROCESSED 4 ITERATIONS

0 ANSWERS

SEARCH TIME: 00.00.01


L3 0 SEA SSS FUL L1

=>

Uploading C:\Program Files\Stnexp\Queries\10714255.str

10714255

6/9/05



```
chain nodes :
1  2  3  4  17  24  25  26  27  28  29  30  31
ring nodes :
5  6  7  8  9  10  11  12  13  14  15  16  18  19  20  21  22  23
chain bonds :
1-2  1-3  1-4  1-24  3-5  4-11  13-17  17-18  24-25  25-26  26-27  26-28  28-29
29-30  30-31
ring bonds :
5-6  5-10  6-7  7-8  8-9  9-10  11-12  11-16  12-13  13-14  14-15  15-16  18-19
18-23  19-20  20-21  21-22  22-23
exact/norm bonds :
1-2  1-3  1-4  3-5  4-11  13-17  17-18  24-25  25-26  26-27  28-29  29-30  30-31
exact bonds :
1-24  26-28
normalized bonds :
5-6  5-10  6-7  7-8  8-9  9-10  11-12  11-16  12-13  13-14  14-15  15-16  18-19
18-23  19-20  20-21  21-22  22-23
```

```
Match level :
1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:CLASS 18:Atom 19:Atom
20:Atom 21:Atom 22:Atom 23:Atom 24:CLASS 25:CLASS 26:CLASS 27:CLASS
28:CLASS 29:CLASS 30:CLASS 31:CLASS
```

L4 STRUCTURE UPLOADED

=> d 14

L4 HAS NO ANSWERS

L4 STR

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

Structure attributes must be viewed using STN Express query preparation.

10714255

6/9/05

=> s l4

SAMPLE SEARCH INITIATED 12:00:28 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 0 TO ITERATE

100.0% PROCESSED 0 ITERATIONS
SEARCH TIME: 00.00.03

0 ANSWERS

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**

PROJECTED ITERATIONS: 0 TO 0
PROJECTED ANSWERS: 0 TO 0

L5 0 SEA SSS SAM L4

=> s l4 ful

FULL SEARCH INITIATED 12:00:36 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 0 TO ITERATE


100.0% PROCESSED 0 ITERATIONS
SEARCH TIME: 00.00.01

0 ANSWERS

L6 0 SEA SSS FUL L4

=>

Uploading C:\Program Files\Stnexp\Queries\10714255.str



chain nodes :

1 2 3 4 17 24 25 26 27 28 29 30 31

ring nodes :

5 6 7 8 9 10 11 12 13 14 15 16 18 19 20 21 22 23

chain bonds :

1-2 1-3 1-4 1-24 3-5 4-11 13-17 17-18 24-25 25-26 26-27 26-28 28-29
29-30 30-31

ring bonds :

5-6 5-10 6-7 7-8 8-9 9-10 11-12 11-16 12-13 13-14 14-15 15-16 18-19
18-23 19-20 20-21 21-22 22-23

exact/norm bonds :

1-2 1-3 1-4 3-5 4-11 13-17 17-18 24-25 25-26 26-27 28-29 29-30 30-31

exact bonds :

1-24 26-28

10714255

6/9/05

normalized bonds :

5-6 5-10 6-7 7-8 8-9 9-10 11-12 11-16 12-13 13-14 14-15 15-16 18-19
18-23 19-20 20-21 21-22 22-23

Match level :

1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:CLASS 18:Atom 19:Atom
20:Atom 21:Atom 22:Atom 23:Atom 24:CLASS 25:CLASS 26:CLASS 27:CLASS
28:CLASS 29:CLASS 30:CLASS 31:CLASS

L7 STRUCTURE UPLOADED

=> s 17

SAMPLE SEARCH INITIATED 12:01:38 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 0 TO ITERATE

100.0% PROCESSED 0 ITERATIONS 0 ANSWERS
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
 BATCH **COMPLETE**
PROJECTED ITERATIONS: 0 TO 0
PROJECTED ANSWERS: 0 TO 0

L8 0 SEA SSS SAM L7

=> s 17 ful

FULL SEARCH INITIATED 12:01:48 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 0 TO ITERATE

100.0% PROCESSED 0 ITERATIONS 0 ANSWERS
SEARCH TIME: 00.00.01

L9 0 SEA SSS FUL L7

=> logoff y

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	485.28	485.49

STN INTERNATIONAL LOGOFF AT 12:01:54 ON 09 JUN 2005

10714255

Iron center, substrate recognition and mechanism of peptide deformylase

Eubacterial proteins are synthesized with a formyl group at the N-terminus which is hydrolytically removed from the nascent chain by the mononuclear iron enzyme peptide deformylase. Catalytic efficiency strongly depends on the identity of the bound metal. We have determined by X-ray crystallography the Fe^{2+} , Ni^{2+} and Zn^{2+} forms of the *Escherichia coli* enzyme and a structure in complex with the reaction product Met-Ala-Ser. The structure of the complex, with the tripeptide bound at the active site, suggests detailed models for the mechanism of substrate recognition and catalysis. Differences of the protein structures due to the identity of the bound metal are extremely small and account only for the observation that Zn^{2+} binds more tightly than Fe^{2+} or Ni^{2+} . The striking loss of catalytic activity of the Zn^{2+} form could be caused by its reluctance to change between tetrahedral and five-fold metal coordination believed to occur during catalysis.

N-terminal formylation and subsequent deformylation appears to be a characteristic feature of eubacterial protein synthesis¹. Peptide deformylase (PDF, EC 3.5.1.31), the enzyme responsible for cleaving the formyl group^{2,3}, is essential for bacterial growth⁴ but seems to be absent in eucaryotic cells (see references cited in ref. 5) which makes it an attractive target for the design of new antibiotics. PDF is a mononuclear iron enzyme⁵⁻⁷ of catalytic efficiency ($k_{\text{cat}}/K_{\text{M}} = 10^5 \text{ M}^{-1}\text{s}^{-1}$) comparable to that of other metalloproteases. Fe^{2+} can be replaced by Ni^{2+} without significant loss of catalytic efficiency, whereas the Zn^{2+} form, prepared from the apoenzyme or by displacement of Fe^{2+} or Ni^{2+} proved virtually inactive⁸.

Previous structural work on the Zn^{2+} form by NMR spectroscopy of the core domain (residues 1–147)⁹ and X-ray crystallography of the full-length protein (residues 1–168) at 2.9 Å resolution⁹ implicates Cys 90, His 132, His 136 and a water molecule as metal ligands. Although the overall fold of PDF differs from other metalloenzymes⁸, the finding that the two histidine ligands reside in the sequence motif HEXXH¹⁰, known^{11,12} to be involved in zinc binding in metalloproteases like thermolysin¹³, has led to the assumption of a similar enzyme mechanism⁹. Recently, we have solved the X-ray structure of the catalytically active Ni^{2+} form at 1.9 Å resolution in complex with the competitive inhibitor PEG identifying the substrate binding site¹⁴. Now we report the structures of the Fe^{2+} , Ni^{2+} and Zn^{2+} forms of the enzyme along with the structure in complex with the reaction product Met-Ala-Ser, and derive detailed models for the mechanism of substrate recognition and catalysis that will help in the design of new antibacterial agents.

Crystal structures

We have solved the crystal structures of the native Fe^{2+} -enzyme in complex with PEG (PDF-Fe/PEG), the nearly inactive Zn^{2+} -forms in the presence (PDF-Zn/PEG) and absence (PDF-Zn) of PEG, the catalytically active Ni^{2+} -form in complex with the product Met-Ala-Ser (PDF-Ni/MAS), and the Zn^{2+} -form cocrystallized with substrate formyl-Met-Ala-Ser (PDF-Zn/fMAS). The structures were solved by molecular

replacement using the PDF-Ni/PEG structure¹⁴ as starting model. Data sets and results of structure determinations are summarized in Table 1 which includes the entries of the previously described PDF-Ni and PDF-Ni/PEG crystals¹⁴ to simplify the discussions.

All crystals possess space group symmetry C2 and contain three molecules in the asymmetric unit⁶, named A, B and C. Cell constants are $a = 140.8 \text{ Å}$, $b = 63.4 \text{ Å}$, $c = 86.8 \text{ Å}$, $\beta = 120.6^\circ$ for PDF/PEG, $a = 143.4 \text{ Å}$, $b = 64.0 \text{ Å}$, $c = 84.6 \text{ Å}$, $\beta = 123.2^\circ$ for PDF, and $a = 143.4 \text{ Å}$, $b = 64.1 \text{ Å}$, $c = 84.9 \text{ Å}$, $\beta = 123.3^\circ$ for PDF/(f)MAS crystals; they do not depend on the type of the bound metal. Typical deviations in bond length, bond angles, dihedral and improper angles from their ideal values in the atomic models are 0.011 Å, 1.2°, 25.5°, and 2.1° respectively. Refined¹⁵ metal occupancies in each structure are at least 0.92 and have been reset to 1.0 in the final structures.

Peptide binding site

Analysis of cocrystals of PDF- Ni^{2+} with Met-Ala-Ser (PDF-Ni/MAS), a product of PDF catalysis, reveals clear electron density for the tripeptide in all three crystallographically independent monomers (Fig. 1a). Co-traces of the three monomers together with bound Ni^{2+} and tripeptide are shown superimposed in Fig. 1b. Generally, the monomer structures are rather similar except for the more flexible regions 62–68 and 148–168 as observed for the PDF-Ni structures¹⁴. More subtle structural differences are found in regions far from direct crystal contacts, notably in the position of water W2. In contrast to all other structures, W2 is found in close proximity of the metal in monomers A and C of the PDF/(f)MAS structures which makes it a fifth ligand (Table 2). Ligation of W2 to the metal is accompanied by a shift of ~0.4 Å of the turn Gly 89–Leu 91 which leads to a more open metal center.

The tripeptide binding region of the PDF-Ni/MAS structure (Fig. 1b) overlaps well the the PEG binding region of the PDF-Ni/PEG structure (Fig. 1 of ref. 14). Since PEG is a competitive inhibitor of PDF activity¹⁴, we can reasonably assume that Met-Ala-Ser binds to the enzyme in a similar manner as formyl-Met-Ala-Ser which is a substrate of the enzyme. The binding of formyl-Met-Ala-Ser can be modeled by shifting Met-Ala-Ser (Fig. 1c) towards the metal by 0.8 Å and replacing W2 by the carbonyl oxygen of the formyl group (Fig. 1d). This hypothetical model of the enzyme–substrate complex, generated using X-PLOR¹⁵ with harmonic restraints to the starting structure, obeys stereochemical constraints and requires only minor changes in the active site region of the protein as observed for the PDF-Ni/MAS structure. Moreover, it explains why PDF has a much lower activity for peptides acetylated at the N-terminus^{2,16,17}. In the model the additional methyl group (which substitutes for the hydrogen in the formyl group) would overlap with the side chains of Leu 91 and Leu 46 and the carbonyl oxygen of Gly 45, as noted by Chan *et al.*⁹. Also, displacement of Leu 91, which in our model of the enzyme mechanism would be part of the oxyanion hole, is likely to reduce catalytic efficiency. Thus, we speculate that substrate recognition by the enzyme may involve most of the interactions between the peptide and PDF as seen in the PDF-Ni/MAS structure.

Interactions of Met-Ala-Ser with PDF are depicted in Fig. 2. The N-terminal amino group of the peptide forms hydrogen bonds with Gly 45, Glu 133 and the waters W1 and W2, whereas the methionine side chain fits neatly into a hydrophobic pocket formed by residues Ile 44, Ile 86, Glu 88, Leu 125, Ile 128, Cys 129 and His 132 (Fig. 2b). Clearly, amino acids like

letters




Fig. 1 **a**, Omit map of Met-Ala-Ser in the PDF-Ni/MAS structure contoured at 1 σ . Figure prepared with BobScript²⁵. **b**, Stereo-view of superimposed C α -traces of the three crystallographically independent copies of peptide deformylase in complex with the reaction product Met-Ala-Ser. The numbers refer to amino acid residues. The transformations for optimal superposition were determined from equivalent C α -atoms using molecule A as a reference and applied to the Ni²⁺ ion (marked as Ni) and the peptide as well. Molecule complexes A, B, C are shown in green, magenta, blue, respectively. **c**, Active site of PDF-Ni (monomer A) with bound catalytic product Met-Ala-Ser, ordered water molecules W1, W2 and the Ni²⁺ ion. **d**, A hypothetical model of PDF with bound substrate formyl-Met-Ala-Ser.

phenylalanine or ethionine instead of methionine also bind to PDF (unpublished results, S.S. and A.F.V.W.) and would also fit into the pocket pointing to significance for inhibitor design. Consistent with the enzyme's low sequence specificity, all hydrogen bonds involve only atoms of the peptide backbone. Moreover, as shown in Figs. 1c,d and 2b, the tripeptide is bound at a position that leaves enough room to accommodate bulky side chains for the second and following residues. Compared with longer substrates or with formyl-Met-amide, the minimal substrate formyl-Met has a much lower catalytic efficiency⁷ which points to the importance of the interaction with Gly 89. Perhaps the presence of the hydrogen bond is communicated to the metal ligand Cys 90 thereby reducing the energy of the transition state. Interestingly, Cys 90 is located in the β -turn Gly 89–Leu 91 which together with water W2 is susceptible to weak crystal packing forces.

Motivated by the finding that PDF-Zn²⁺ is nearly catalytically inactive, we have determined its structure using crystals grown in the presence of the substrate formyl-Met-Ala-Ser (data set PDF-Zn/fMAS). Unfortunately, because of residual activity of the Zn²⁺ form in the crystal or because of small contamination by more active metallo forms of the enzyme, no density was found in the map that could be assigned to the formyl group; this is in sharp contrast to the clear density seen for the peptide product Met-Ala-Ser. In fact, both the Ni²⁺- and Zn²⁺-forms of PDF with the bound tripeptide are remarkably similar.

Metal center

The various metal forms of the PDF/PEG, PDF and PDF/(f)MAS crystals, each grown under nearly identical chemical conditions, allow us to study the influence of the metal on the structure. In agreement with earlier




Fig. 2 Peptide deformylase in complex with the reaction product Met-Ala-Ser. **a**, Peptide binding scheme; Gly 45, Glu 88, Gly 89, His 132 and Glu 133 are conserved residues. Dashed lines indicate hydrogen bonds. Mean distances between donor and acceptor atoms as observed in the three crystallographically independent monomers are given in Å. The distance between the N-terminal amino group of the peptide to the Ni^{2+} -ion is 3.9 Å. **b**, Protein atoms (white, carbon; red, oxygen; dark blue, nitrogen; green, sulphur) and catalytic metal (magenta, Ni^{2+}) are depicted as space-filling spheres. Met-Ala-Ser is in ball-and-stick representation with carbon atoms and bonds colored yellow. Water molecules W1, W2 are shown as small light blue spheres. Figure prepared using MOLSCRIPT²⁶ and Raster3D²⁷.

NMR³ and X-ray studies of PDF-Zn⁹, the metal ion in the structures presented here (except those of the peptide complexes) is tetrahedrally ligated. Ligands are the Sγ-atom of Cys 90, the Ne2-atoms of His 132 and His 136 in the HEXXH motif, and an oxygen atom of the group W1 modeled as a water molecule. All ligands, occupying well defined positions in the electron density maps, are precisely aligned by an intricate network of hydrogen bonds involving conserved residues of the enzyme family⁴. As shown in Table 2, the arrangement of the ligands, in particular of W1, deviates significantly from ideal tetrahedral symmetry in all structures.

Considering the variability found in the metal ligand positions for the three crystallographically independent molecules in each structure, it appears that crystal packing effects or random errors in the atomic models obscure all structural differences between the Fe^{2+} , Ni^{2+} and Zn^{2+} forms of the enzyme. In fact, the influence of the metal on the structure is extremely small and barely detectable. It is known that Zn^{2+} is bound more tightly by the enzyme than the other metals⁶. This effect becomes visible in the structures if the tetrahedral volumes, spanned by the four metal ligands, are compared as shown in Table 2. In all cases, volumes around the Zn^{2+} ion are smaller than in the presence of Fe^{2+} or Ni^{2+} .

The extreme similarity among the structures in each of the three groups of crystals has also been confirmed by Fourier analysis of differences between observed structure factors.

This analysis is known to be very sensitive, yet robust towards errors in the atomic model. Moreover, we have excluded the metal from calculating model phases. In all cases, the dominant maxima in the resulting difference maps varied between 10σ and 36σ and account for the difference of two or four electrons between the compared metal forms. Minima and minor maxima are predominantly found in the vicinity of the ligand Cys 90. These structural differences are remarkably small and of a magnitude well below the contribution expected from an ordered water molecule that may be present, absent or displaced due to the metal type. We conclude that the enormous differences in catalytic activity of the metal forms of the enzyme ultimately result from the tighter binding of the Zn^{2+} ion as compared to Fe^{2+} or Ni^{2+} .

Catalytic model

It has been noted that the active site of PDF is similar to thermolysin although its overall fold differs from other metalloproteases⁸. Both proteins use two histidines from the common HEXXH sequence motif for metal ligation and require the glutamate residue for catalytic activity. In fact, mutations of Glu 133 to Ala, Asp or Gln were found to eliminate catalytic activity of PDF^{18,19}. This has led to the proposal of a mechanism of deformylation⁹ that resembles the mechanism suggested for thermolysin²⁰. Below we propose a model of the catalytic cycle of PDF that is consistent with the available

letters




Fig. 3 Proposed reaction cycle of peptide deformylase.

structures and biochemical data. Besides being more detailed, it differs from the model of Chan *et al.*⁹ by the inclusion of water W2 and the role of residues Leu 91 and Gln 50 which form an oxyanion hole as described for serine and cysteine proteases²¹.

The reaction cycle could proceed in the following steps (Fig. 3). Step 1 represents the initial state of PDF as found in the PDF-Ni and PDF-Zn structures. In the next step the formylated peptide binds as shown in the hypothetical model of the enzyme-substrate complex (Fig. 1d) thereby replacing water W2 by the carbonyl oxygen of the formyl group. The carbonyl oxygen is polarized by hydrogen bonds to the amide of Leu 91 and the side chain of Gln 50. This supports the nucleophilic attack of W1 (probably a deprotonated water) on the carbonyl carbon of the formyl group that leads to the transition state of the reaction as depicted in step 3. The carbonyl oxygen is tetrahedrally ligated by the metal, carbonyl carbon, side chain amide of Gln 50, and main chain amide of Leu 91. This arrangement of amide hydrogen bond donors resembles the oxyanion hole in serine and cysteine proteases for stabilizing the transition state²¹. The absence of ionizable groups in the oxyanion hole is well in line with the broad pH-activity profile of PDF-Fe which is nearly constant between pH 6.1 and 11.2 (ref. 7). The electronic state of the attacked carbonyl carbon changes from sp^2 to sp^3 accompanied by a transition from the

tetrahedral to a five-coordinate metal center, this transition being strongly inhibited in the case of PDF-Zn due to the tighter binding of this metal ion as compared with Fe^{2+} or Ni^{2+} . Probably the rearrangement of the metal ligands is supported indirectly by the hydrogen bond between Gly 89 and the peptide substrate as mentioned above.

The proton of W1 is transferred with the help of Glu 133 to the amide at the N-terminus of the peptide, the added positive charge making the nitrogen suitable as a leaving group. Subsequent bond cleavage leads to the ternary enzyme-formate-peptide complex shown in step 4. Here, the formate is bound to the five-coordinated metal center and the free N-terminus of the peptide is hydrogen bonded to Glu 133 as found in the PDF-Ni/MAS structure. The reaction proceeds by dissociation of the peptide which leaves an activated enzyme-formate complex as shown in step 5. The existence of such a complex is consistent with biochemical data which show that PDF can transfer the formyl group from one formyl peptide to another peptide by a 'ping-pong' mechanism⁷. The reaction cycle closes with the release of formate and the uptake of two water molecules, W1 and W2. It is very likely that formate release proceeds by attack of a water molecule on the metal rather than on the formate carbon atom. Evidence comes from the lack of ^{17}O exchange during incubation of ^{17}O formate in the presence of PDF⁷.

Table 1 Data collection and refinement statistics

Compound	PDF-Fe/PEG	PDF-Ni/PEG	PDF-Zn/PEG	PDF-Ni	PDF-Zn	PDF-Ni/MAS	PDF-Zn/fMAS
PDB accession number	1BSZ	1ICJ	1BS4	1BS7	1BS5	1BS6	1BS8
Outer shell (Å)	2.01–1.9	2.01–1.9	2.01–1.9	2.64–2.5	2.64–2.5	2.23–2.1	2.33–2.2
Measured reflections	104,453	281,561	188,577	90,069	130,772	118,387	117,576
Unique reflections	50,232	51,138	51,676	21,929	22,399	37,363	32,674
Completeness (%)	96.1	97.9	99.2	97.2	99.3	98.0	99.2
R _{sym} (%) ¹	5.9	7.5	4.8	8.6	7.6	5.6	8.0
<I/σ> in outer shell	2.2	2.7	2.8	2.4	1.9	1.8	2.0
R-factor (%) ²	19.7	19.3	19.3	20.3	20.8	20.6	20.4
R _{free} (%) ³	24.7	23.8	24.0	27.2	25.8	25.8	26.1
Coordinate error (Å)	0.23	0.22	0.22	0.33	0.33	0.29	0.29
Protein atoms	4,122	4,122	4,122	4,038	4,038	4,107	4,107
Alternate locations	84	84	84	0	0	69	69
Heterogen atoms	97	97	97	13	13	73	73
Solvent molecules	211	211	211	132	132	165	165

¹R_{sym} = $\sum_h \sum_i |I_{hi} - I_h| / \sum_h I_h$ where h are unique reflection indices and I_{hi} are the intensities of symmetry equivalent reflections giving a mean value of I_h .

²R = $\sum |F_{obs} - F_{model}| / \sum F_{obs}$ where F_{obs} and F_{model} are observed and atomic model structure factor amplitudes respectively.

³R-factor calculated for 10% of randomly chosen reflections that were excluded from the refinement

Methods

Preparation of all crystals except PDF-Zn/fMAS, enzymatic activity tests and determinations of metal content were performed as described⁶. PDF-Zn/fMAS crystals were grown under conditions similar to those for PDF-Ni/MAS with 20 mM formyl-Met-Ala-Ser instead of MAS and in addition 500 mM ammonium formate and 100 mM KF.

Diffraction data were collected at room temperature by the rotation method and recorded by an electronic area detector (X-rays: CuKα, focused by Franks double-mirror optics; generator: GX-18, Elliot/Enraf-

Nonius, Delft, operated at 35 kV, 50 mA; detector: X100, Siemens/Nicolet; crystal to detector distance: 10 cm; rotation/image: 0.0417° or 0.0833°). Integrated intensities were extracted from the rotation images by the program package XDS²² which includes routines for scaling data from several crystals as well as for space group determination from the observed diffraction pattern²³.

All structures were solved by molecular replacement using the PDF-Ni/PEG structure¹⁴ as starting model, followed by model correction²⁴ and refinement¹⁵. For comparison of two isomorphous crys-

Table 2 Metal coordination¹

Compound	Bond angles ² (°)						Bond length ³ (Å)					V ⁴ (Å ³)
	α ₁₂	α ₁₃	α ₁₄	α ₂₃	α ₂₄	α ₃₄	d ₁	d ₂	d ₃	d ₄	d ₅ ⁵	
PDF-Fe/PEG	119	105	128	103	103	93	2.3	2.1	2.1	2.2	3.7	5.0
	108	112	131	101	107	95	2.3	2.3	2.1	2.0	3.7	5.0
	116	111	132	103	104	85	2.2	2.2	2.2	2.3	3.6	5.1
PDF-Ni/PEG	116	102	133	100	98	103	2.2	2.1	2.1	1.9	3.7	4.2
	111	105	134	104	100	98	2.3	2.2	2.0	2.0	3.6	4.6
	115	104	133	102	102	95	2.3	2.1	2.1	2.1	3.7	4.7
PDF-Zn/PEG	118	102	125	104	101	104	2.3	2.1	2.0	1.8	3.9	4.1
	117	105	124	104	100	103	2.3	2.0	2.1	1.9	3.8	4.3
	116	103	126	106	101	102	2.2	2.1	2.1	2.0	3.8	4.6
PDF-Ni	127	107	137	96	95	73	2.1	2.1	2.4	2.5	3.9	4.8
	115	116	137	107	92	84	2.2	2.2	2.0	1.9	3.7	4.0
	114	110	149	99	92	79	2.2	2.2	2.1	2.2	n.o.	4.2
PDF-Zn	118	102	131	107	101	93	2.1	2.0	2.2	2.2	4.1	4.7
	118	107	127	109	96	96	2.1	2.1	2.1	1.8	3.7	4.0
	115	107	143	105	90	91	2.1	2.1	2.1	1.8	n.o.	3.8
PDF-Ni/MAS	121	105	157	98	81	77	2.4	2.2	2.3	2.2	2.3	4.3
	122	111	142	100	88	84	2.3	2.2	2.2	1.9	3.4	4.3
	115	98	155	102	89	82	2.4	2.2	2.2	2.0	2.7	4.2
PDF-Zn/fMAS	117	109	156	101	83	77	2.1	2.3	2.1	1.9	2.3	3.6
	116	104	133	113	93	96	2.2	2.1	2.0	1.9	3.7	4.1
	115	109	154	102	87	77	2.2	2.0	2.0	2.3	2.2	3.8

¹Bond angles, bond lengths and volumes are given for monomers A, B, C in the asymmetric unit of each compound.

²Bond angle α_{ij} is the angle between ligand i, metal and ligand j (i, j = 1, ..., 4).

³Bond lengths d₁, d₂, d₃, d₄ are distances from the central metal to its four ligands, Cys 90 (Sγ), His 132 (Nε2), His 136 (Nε2), and water molecule W1 respectively.

⁴V is the volume of the tetrahedron formed by the four metal ligands.

⁵d₅ is the distance between water molecule W2 and the metal. W2 is not observed in monomer C of the PDF-Ni and PDF-Zn structures.

letters

tal structures difference electron density maps were computed from structure factors calculated as difference between the measured amplitudes of each data set multiplied by a phase factor derived from the atomic model of one of the crystals.

Coordinates. Protein Data Bank accession numbers are given in Table 1.

Acknowledgments

We thank D. Madden, K. Fritz-Wolf and K. Scheffzek for critical discussions and help at various stages of the project, H. Wagner for excellent maintenance of the X-ray facilities at the MPI Heidelberg, I. Dehof for help with the figures, and K. Holmes and J. Knappe for continuous support.

Andreas Becker¹, Ilme Schlichting², Wolfgang Kabsch¹, Dieter Groche³, Sabine Schultz³ and A. F. Volker Wagner³

¹Max-Planck-Institut für medizinische Forschung, Abteilung Biophysik, Jahnstraße 29, 69120 Heidelberg, Germany. ²Max-Planck-Institut für molekulare Physiologie, Abteilung Physikalische Biochemie, Rheinlanddamm 201, 44139 Dortmund, Germany. ³Biochemie-Zentrum Heidelberg, Ruprecht-Karls Universität, Im Neuenheimer Feld 501, 69120 Heidelberg, Germany.

Correspondence should be addressed to W.K. email: kabsch@mpimf-heidelberg.mpg.de

Received 22 June, 1998; accepted 11 September, 1998.

1. Kozak, M. *Microbiol. Rev.* **47**, 1–45 (1983).
2. Adams, J.M. *J. Mol. Biol.* **33**, 571–589 (1968).
3. Ball, L. A. & Kaesberg, P. *J. Mol. Biol.* **79**, 531–537 (1973).
4. Mazel, D., Pochet, S. & Marlière, P. *EMBO J.* **13**, 914–923 (1994).
5. Rajagopalan, P.T.R., Yu, X.C. & Pei, D. *J. Am. Chem. Soc.* **119**, 12418–12419 (1997).
6. Groche, D. et al. *Biochem. Biophys. Res. Commun.* **246**, 342–346 (1998).
7. Groche, D. Ph.D. thesis, Universität Heidelberg. Charakterisierung des Eisenzentrums und des Katalysenmechanismus von Peptid-Deformylase aus *Escherichia coli* (1995).
8. Meinzel, T., Blanquet, S. & Dardel, F. *J. Mol. Biol.* **262**, 375–386 (1996).
9. Chan, M.K. et al. *Biochemistry* **36**, 13904–13909 (1997).
10. Meinzel, T. & Blanquet, S. *J. Bacteriol.* **175**, 7737–7740 (1993).
11. Vallee, B.L. & Auld, D.S. *Biochemistry* **29**, 5647–5659 (1990).
12. Jongeneel, C.V., Bouvier, J. & Balroch, A. *FEBS Lett.* **242**, 211–214 (1989).
13. Holmes, M.A. & Matthews, B.W. *J. Mol. Biol.* **160**, 623–629 (1982).
14. Becker, A., Schlichting, I., Kabsch, W., Schultz, S. & Wagner, A.F.V. *J. Biol. Chem.* **273**, 11413–11416 (1998).
15. Brünger, A.T. *J. Mol. Biol.* **203**, 803–816 (1988).
16. Wei, Y. & Pei, D. *Anal. Biochem.* **250**, 29–34 (1997).
17. Meinzel, T. & Blanquet, S. *J. Bacteriol.* **177**, 1883–1887 (1995).
18. Meinzel, T., Lazennec, C. & Blanquet, S. *J. Mol. Biol.* **254**, 175–183 (1995).
19. Meinzel, T., Lazennec, C., Villioing, S. & Blanquet, S. *J. Mol. Biol.* **267**, 749–761 (1997).
20. Matthews, B.W. *Acc. Chem. Res.* **21**, 333–340 (1988).
21. Ménard, R. & Storer, A.C. *Biol. Chem. Hoppe-Seyler* **373**, 393–400 (1992).
22. Kabsch, W. *J. Appl. Crystallogr.* **21**, 916–924 (1988).
23. Kabsch, W. *J. Appl. Crystallogr.* **26**, 795–800 (1993).
24. Jones, T.A., Zou, J.Y., Cowan, S.W. & Kjeldgaard, M. *Acta crystallogr. A* **47**, 110–119 (1991).
25. Esnouf, R.M. *J. Mol. Graphics* **15**, 132–134 (1997).
26. Kraulis, P.J. *J. Appl. Crystallogr.* **24**, 946–950 (1991).
27. Merritt, E.A. & Murphy, M.E.P. *Acta Crystallogr. D* **50**, 869–873 (1994).

MutY catalytic core, mutant and bound adenine structures define specificity for DNA repair enzyme superfamily

The DNA glycosylase MutY, which is a member of the Helix-hairpin-Helix (HhH) DNA glycosylase superfamily, excises adenine from mispairs with 8-oxoguanine and guanine. High-resolution crystal structures of the MutY catalytic core (cMutY), the complex with bound adenine, and designed mutants reveal the basis for adenine specificity and glycosyl bond cleavage chemistry. The two cMutY helical domains form a positively-charged groove with the adenine-specific pocket at their interface. The Watson-Crick hydrogen bond partners of the bound adenine are substituted by protein atoms, confirming a nucleotide flipping mechanism, and supporting a specific DNA binding orientation by MutY and structurally related DNA glycosylases.

Reactive oxygen species that are generated during aerobic respiration and immune response cause oxidative lesions in DNA that are repaired by the base excision repair pathway¹. Oxidative DNA damage is implicated in mutagenesis, carcinogenesis and aging². Hydroxyl radicals rapidly react with the guanine C8 position producing a steady-state level of mutagenic 8-oxoguanines (8OG) in cells³. Oxidation also creates 8-oxo-GTP that, like 8OG in template DNA, is generally mispaired with adenine by replicative polymerases, creating A-T to C-G and G-C to T-A transversion mutations⁴. In *Escherichia coli*, three key enzymes, MutT, MutM (FPG), and MutY, function synergistically to protect cells from the deleterious effects of guanine oxidation^{5,6}. MutT removes 8-oxo-GTP⁷ to reduce its misincorporation by polymerases, and MutM excises 8OG from

8OG-C DNA base pairs⁸. Only the mismatched-adenine specific glycosylase MutY recognizes the mutational intermediate 8OG-A mispair⁹, and loss of the *E. coli mutY* gene leads to elevated mutation rates⁸. Mutations in the three analogous human enzymes may increase cancer susceptibility due to elevated transversion frequencies^{6,9}. Importantly, among these three enzymes, only the sequence of MutY is conserved from bacteria to humans suggesting its fundamental importance in DNA repair¹⁰. Moreover, as only the 8OG in 8OG-A mispairs is correctly coding, restoration of the original G-C base pair requires a two step process. First, MutY excises adenine mispaired with 8OG, creating an abasic site that is processed by the multi-enzyme base excision repair pathway^{5,11} to yield an 8OG-C base pair. Second, MutM removes 8OG to initiate a final cycle of base excision repair that restores the original G-C base pair.

Cloning and sequencing of *mutY*¹² and other DNA repair genes revealed a base excision repair glycosylase superfamily¹³, which includes both pure DNA glycosylases, like 3-methyladenine glycosylase II (AlkA), and DNA glycosylase/lyases, like endonuclease III (EndoIII). Crystal structures of EndoIII¹⁴ and AlkA^{15,16} show that the most conserved superfamily structural element is the Helix-hairpin-Helix (HhH) motif¹⁷. However, no crystal structures with bound DNA or with bound target bases exist for any member of the HhH glycosylase superfamily. Thus, the structural and chemical basis of DNA binding, substrate specificity and glycosyl bond cleavage by MutY and other members of the HhH DNA glycosylase superfamily is largely uncharacterized.

MutY structure

Here we characterize the structural chemistry of MutY and HhH glycosylase catalysis and recognition of damaged bases using: three designed MutY mutants, MutY inhibition by relevant base analogs, and high-resolution crystal structures of the fully-active, 26,000 M_r catalytic core of MutY (cMutY) that lacks the 13,000 M_r C-terminal region of unknown function¹⁸. Three cMutY structures were determined: wild type cMutY (1.4 Å resolution), an inactive site-directed

Copyright © 2003 EBSCO Publishing